

Comment on Liu et al: ‘Final state interactions in the decays $J/\Psi \rightarrow VPP$ ’

D. V. Bugg¹

Queen Mary, University of London, London E1 4NS, UK

Abstract

Statements concerning Extended Unitarity (EU) are clarified.

PACS numbers: 11.80.Et, 1360.6e, 14.40.Cs.

The objective of this Comment is to expose points of uncertainty and disagreement concerning Extended Unitarity (EU). Liu et al. [1] make several assertions criticising an earlier publication of mine [2], where fits to four sets of data are compared with the form of Extended Unitarity formulated by Au, Morgan and Pennington [3]. The principles should apply to the overlap of any two resonances having the same quantum numbers. Both my publication and that of Liu et al. concern σ and $f_0(980)$ where they overlap near 1 GeV. Here σ is a shorthand for the broad $\pi\pi$ component going through a phase shift of 90° in the mass range 900–940 MeV.

For $\pi\pi$ elastic scattering, a Breit-Wigner resonance has an elastic scattering amplitude

$$\rho_{\pi\pi} T_{\pi\pi \rightarrow \pi\pi} = \frac{M\Gamma_{\pi\pi}(s)}{M^2 - s - m(s) - iM\Gamma_{tot}(s)} = \frac{N(s)}{D(s)} \quad (1)$$

$$m(s) = \frac{1}{\pi} P \int \frac{M\Gamma_{tot}(s') ds'}{s' - s}; \quad (2)$$

ρ is $\pi\pi$ phase space and $N(s)$ is real. Watson’s theorem [4] amounts to the fact that the Breit-Wigner denominator $D(s)$ and the associated pole are universal. The numerator is not universal.

Fig. 1(a) shows the $\pi\pi$ mass projection from BES II data for $J/\Psi \rightarrow \omega\pi^+\pi^-$ [5]. The peak near 500 MeV may be fitted with the σ pole taking $N(s) = 1$, unlike elastic scattering where $N(s)$ is constrained by unitarity to be $M\Gamma_{\pi\pi}(s)$. There may be a small dip in the data due to $f_0(980)$ in the bin just above the KK threshold. The σ amplitude has a different s -dependence to elastic scattering and the relative magnitude of $f_0(980)$ to σ is also different.

Let us denote the $\pi\pi$ production amplitude by $A_k(s)$, where k refers to channels $\pi\pi$, KK , etc. Liu et al. use the relation

$$\text{Im}(A_i) = -\rho_k T_{ik}^* A_k, \quad (3)$$

where T_{ik} is the $2 \rightarrow 2$ amplitude. The form of EU used by Liu et al. is that the phase of T is identical to that of elastic scattering. They do not specify if this is the phase shift δ or the phase ϕ measured from the bottom of the unitarity circle. These differ above the inelastic threshold. The Omnès relation [6] connects the magnitude of the elastic scattering amplitude to ϕ using analyticity, so ϕ appears to be the most relevant angle and will be used here.

The *critical* assumption of EU is that the relative phases of σ and f_0 in T must be identical in production reactions and elastic scattering. This is an assumption going beyond Watson’s theorem [4].

In elastic scattering, σ and f_0 amplitudes are constrained below the KK threshold to move around the unitary circle on the Argand diagram. In Cern-Munich data [7], it is obvious by eye

¹email: david.bugg@stfc.ac.uk




Figure 1: (a) The $\pi\pi$ mass projection for BES II data on $J/\Psi \rightarrow \omega\pi^+\pi^-$; the upper histogram shows the fit to EU in the form of AMP, and the lower histogram the $\pi\pi$ S-wave intensity; (b) an enlarged view of the corner of the Dalitz plot; cross-hairs show the point of intersection of $f_0(980)$ with the strong $b_1(1235)$ component.

that phase shifts of σ and $f_0(980)$ add to a good approximation near 1 GeV. This additivity of phases may be accommodated by multiplying S -matrices $\eta \exp(2i\delta)$, though it is an open question whether η parameters should multiply. For present considerations concerning the overlap of σ and f_0 , this is not critical, since the inelasticity of the σ rises fairly slowly at the KK threshold because the σ has a large $\pi\pi$ width.

For a production process the relative magnitudes of σ and f_0 are different, because of different matrix elements connecting initial and final states. It is not obvious that T of different resonances must combine in the same way as for elastic scattering. In a $2 \rightarrow 2$ process, σ and f_0 amplitudes combine to make an overall amplitude T which obeys the relation

$$\text{Im } T = TT^*. \quad (4)$$

In a $1 \rightarrow 3$ or $2 \rightarrow 3$ process, the boundary condition for the initial state is different. Why can $f_0(980)$ not be completely absent in some production processes? In that case, Eq (4) is still valid for the σ amplitude *alone*, but EU demands that the phase of $f_0(980)$ still appears in the production process despite its magnitude being absent. Note that the process $J/\Psi \rightarrow \omega\pi^+\pi^-$ accounts for only 1% of all J/Ψ decays.

These questions were much discussed in the early 1960s and led to the isobar model, where each pole is multiplied by complex coupling constants and amplitudes are added, not multiplied. For experimentalists, this is a simple form with which to parametrise data, except possibly in the small region where they overlap and EU could play a role. Even there it parametrises average phase differences between resonances. In the production process, the matrix element involves an unknown integral between the initial and final-state wave functions. This is what Liu et al. seek to model. There is no dispute that multiple scattering processes do affect phases.

The Omnès relation, based on analyticity, allows the magnitude of the $\pi\pi$ amplitude to be derived from phases ϕ . A subtlety is that the magnitude can be multiplied by a polynomial. This polynomial accommodates, for example, a form factor arising from the overlap integral between initial and final states. Liu et al. note this polynomial, but do not give details of whether or how they use it.

On Fig. 1(a). the magnitude of the σ amplitude is not the same as for elastic scattering. My approach, consistent with the data within errors, is to take $N(s) = 1$. Liu et al. may use instead an explicit polynomial relating elastic scattering and production, but they do not say. They derive relative magnitudes of σ and f_0 from equations given in earlier work by Lähde and Meissner [8]. This takes σ and f_0 to be $\bar{n}n$ and $\bar{s}s$ linear combinations, though results would be similar assuming 4-quark compositions (apart from SU3 Clebsch-Gordan coefficients). The matter of different SU3 Clebsch-Gordan coefficients is not presently a critical issue for EU, though it would bear upon the relation of σ and f_0 to 2-quark, 4-quark and/or meson-meson components in wave functions. If the relative magnitudes of $f_0(980)$ and σ are the same in $J/\Psi \rightarrow \omega\pi\pi$ data and elastic scattering, the polynomial is 1 (except for standard form factors), but if the relative magnitudes are different, a polynomial is required peaking at the mass of $f_0(980)$ and reproducing its shape. What does this polynomial mean?

From the point of view of the isobar model, this is inconvenient. The idea of the isobar model is that form factors should have small and slowly varying effects, usually negligible. Liu et al do not show the Argand diagram for their amplitude. It needs to be exposed what polynomial $P(s)$ is being used, so the reader knows what price is being paid to preserve EU.

Paradoxes can arise in limiting cases. Suppose the f_0 amplitude is strictly zero in a production process. One then sees the σ amplitude alone. According to Watson's theorem, the production amplitude must then have the phase of the σ . The hypothesis of EU requires a polynomial which exactly cancels the magnitude *and* phase of the f_0 in elastic scattering; but the cancellation of the phase is not allowed, therefore a zero f_0 amplitude appears to be forbidden by EU. My opinion is that this is a critical defect in the hypothesis of EU.

Let us return to details of $J/\Psi \rightarrow \omega\pi\pi$ data. There were two fits reported in the BES publication [5]. The second does contain a large $f_0(980)$ signal. However, this gives a fit worse in χ^2 by over 200 compared to a free fit to $f_0(980)$. For fine details, the reader should consult Ref. [2], but Fig. 1 presents essential points. Fig. 1(b) shows an enlargement of the corner of the Dalitz plot where the f_0 crosses the strong $b_1(1235)$ band in the data; the intersection is shown by the cross-hairs. If there were a strong f_0 component with the same phase as σ , one should see the interference dip between them as a diagonal band running at 45° from top left to bottom right. It should show a distinctive interference as it crosses the $b_1(1235)$ band centred on the vertical cross-hair. This interference should have a full width at half-maximum of 0.07 GeV^2 in $m^2(\omega\pi)$. There is no sign of this interference with $b_1(1235)$, σ or $f_2(1270)$. This is why the fitted $f_0(980)$ components is very small in my fit to the data, Ref. [2].

Liu et al. do not have access to the Dalitz plot. Their Fig. 5 shows structure just below 1 GeV which seems to require an f_0 at least as large as the σ amplitude in this mass region. Firstly it is not obvious how to reconcile this with $J/\Psi \rightarrow \phi\pi^+\pi^-$ data where the σ amplitude is much smaller than the strong $f_0(980)$ peak. Secondly the precipitous drop in the amplitude fitted to $\omega\pi^+\pi^-$ data just below the KK threshold should generate a cliff-like diagonal structure on Fig. 1(b). The fit of Liu et al. is inconsistent with the isobar model fit shown as the dashed curve on Fig. 1(a). The broad diagonal band of Fig. 1(b) arises from interferences of σ with $b_1(1235)$ and $f_2(1270)$.

It would help the discussion if the BES II collaboration would make publicly available the data set produced by myself and collaborators, the corresponding Monte Carlo data and the sidebin events used to subtract the 14% experimental background. The decay plane of the ω provides delicate information defining accurately the $b_1(1235)\pi$ signal, which then serves as an




Figure 2: (a) AFS data compared with the isobar model fit; (b) enlargement near 1 GeV: the full curve is the isobar model fit, the chain curve shows what appears in elastic $\pi\pi$ scattering, and the dashed curve shows the isobar model fit supplemented by a freely fitted $I = 2$ $\pi\pi$ amplitude.

interferometer determining magnitudes and phases of amplitudes crossing it.

A much clearer test of EU arises in data from the ISR [9] for central production of $\pi\pi$ in the process $pp \rightarrow pp(\pi\pi)$. The central $\pi\pi$ pair is far removed in rapidity from final state protons, and there is no evidence for a Deck effect, i.e. resonant $p\pi$ or $p\pi\pi$ combinations in the central region. In this case, EU predicts that relative phases of σ and f_0 should be identical to $2 \rightarrow 2$ processes. The data are easily fitted by the isobar model. It requires relative intensities $f_0(980)/\sigma \sim 60\%$ of elastic scattering at the KK threshold. The fitted f_0 phase is $(57 \pm 7)^\circ$ below the EU prediction. Features of the fit are displayed in Figs. 9 and 10 of Ref. [2]. Fig. 2 summarises essential points.

Fig. 2(a) shows the isobar model fit (which includes a small slowly varying form factor to reproduce the precise shape of the sigma pole up to 0.8 GeV). Fig. 2(b) shows details of the mass range near 1 GeV. The full curve is the isobar model fit without any $I = 2$ $\pi\pi$ amplitude. The chain curve shows what is observed in elastic $\pi\pi$ scattering, suitably normalised. These two curves are very different in the region of $f_0(980)$, hence the phase difference required in the isobar model fit. There could be an $I = 2$ production amplitude. The dashed curve shows the result of fitting this freely. The fit is still poor and the $I = 2$ amplitude is twice as large as in elastic scattering. My view is that these data rule out EU in the form of Liu et al.

Morgan and Pennington [10] did manage to fit central production data with EU but required an additional third-sheet pole at $M = 978 - i28$ MeV. Since then, data on $f_0(980)$ have improved, and there is no sign of this additional narrow third-sheet pole in BES II data.

In Ref. [2], an empirical relation is presented which does fit the ISR data. The assumption is made that the production process can be considered as a black box. Then it is postulated that whatever linear combination of σ and f_0 is produced will rescatter according to the usual relation $Im T = TT^*$, i.e. the pions rescatter as an isolated pair with magnitudes different from elastic scattering. If the asymptotic combination is written $T = T_\sigma + \beta T_{f_0} \exp^{2i\Psi}$ within the elastic regime, and β is real, it can be shown that

$$2\Psi = 2\delta_\sigma - \delta_{f_0} + \sin^{-1}(\beta \sin \delta_{f_0}). \quad (5)$$

If $\beta \neq 1$, this is a different relation from the $2 \rightarrow 2$ process. As β decreases from 1, Ψ falls rapidly. In Ref. [2], the exact form $Im T = TT^*$ is used, including effects of coupling to KK . Although Liu et al. dismiss this approach, it does have the merit of correctly predicting the observed phase difference between σ and f_0 within errors.

No analyses so far include the dispersive cusp at the KK threshold in both σ and $f_0(980)$. The cusp originates from a discontinuity in slope of the KK amplitude at threshold. This causes a corresponding discontinuity in slope of the real part of the amplitude, hence the cusp [11]. I have made some preliminary fits to BES II data including this cusp, and can offer some cautionary comments to others who may try something similar. In order to make the dispersion integral of Eq. (2) converge, a form factor must be included in the KK channel. Whatever form factor is adopted, there is a narrow peak in the real part of the $\pi\pi$ amplitude exactly at the KK threshold. The narrow width implies that mass resolution is of critical importance in any comparison with data. The width of the cusp is quite stable. However, its height above zero real part does depend on exactly what form factor is used. Either this form factor must be predicted from a model of $f_0(980)$ or the height of the cusp must be treated as a variable in fitting data. This source of uncertainty adds to difficulties in deciding the nature of $f_0(980)$.

The advice to experimentalists is to use the isobar model, which is simple and well defined. It takes account empirically of phases such as Liu et al. are trying to fit in a different way. If mass resolution is very good, it will be necessary to include some s -dependence in $\Gamma_{\pi\pi}$ similar to the cusp. Theorists can then compare the fit with models of f_0 and the cusp those models predict. It may well be necessary to allow for $f_0 \rightarrow \eta\eta$.

In conclusion, the free fit made by Liu et al. to intermediate steps in $J/\Psi \rightarrow \omega\pi^+\pi^-$ accounts for phases used in the isobar model, but the magnitude of $f_0(980)$ signal they fit looks inconsistent with Fig.1(b). The ISR data are in strong disagreement with EU. In those data, there should be no significant perturbations due to other resonances and the production mechanism is clean.

References

- [1] B. Liu et al., arXiv: 0901.1185
- [2] D.V. Bugg, Eur. Phys. J C54 (2008) 73 [arXiv: 0801.1908]
- [3] K.L. Au, D. Morgan and M.R. Pennington, Phys. Rev. D35 (1987) 1633
- [4] K.M. Watson, Phys. Rev. 88 (1952) 1173
- [5] M. Ablikim et al. (BES II Collaboration) Phys. Lett. B598 (2004) 149 [arXiv: hep-ex/0406038]
- [6] R. Omnès, Nu. Cim. 8 (1958) 316
- [7] B. Hyams et al., Nucl. Phys. B64 (1973) 134
- [8] T.A. Lähde and U.-G. Meissner, Phys. Rev. D74 (2006) 034021 [arXiv: hep-ph/0606133]
- [9] T. Åkesson et al., Nucl. Phys. B264 (1986) 154
- [10] D. Morgan and M.R. Pennington, Phys. Rev. D48 (1993) 11851
- [11] D.V. Bugg, J. Phys. G: Nucl. Phys. 35 (2008) 075005 [arXiv: 0802.0934]